

## Cardinal invariants about shrinkability of unbounded sets

Masaru Kada<sup>a,\*</sup>, Yoshifumi Yuasa<sup>b,1</sup>

<sup>a</sup> University of Osaka Prefecture, Sakai, Osaka 593, Japan

<sup>b</sup> Waseda University, Shinjuku, Tokyo 169, Japan

Received 17 July 1995; revised 2 October 1995

### Abstract

In our previous paper (Eda et al., to appear), we introduced a cardinal invariant  $b^*$  and studied some properties of the cardinal  $b^*$ . In the present paper we define new cardinal invariants which are related to Cichoń's diagram and generalize the notion of  $b^*$ . We investigate the relations between them and other cardinals which appear in Cichoń's diagram.

**Keywords:** Cardinal invariants; Measure; Category

**AMS classification:** 03E05; 03E35

### 1. Introduction

In our previous paper [5], we introduced a cardinal invariant  $b^*$ , which is defined with the notion of unbounded family in  $\omega^\omega$  but differs from  $b$ . It was a problem in set-theoretic topology that motivated us to consider such a new cardinal invariant.

Let  $\kappa$  be an infinite cardinal. The *sequential fan*  $S_\kappa$  is the following topological space:  $S_\kappa = \{\infty\} \cup (\kappa \times \omega)$  as a set, every point of  $\kappa \times \omega$  is isolated, and a basic neighborhood of  $\infty$  is of the form  $U_\varphi = \{\infty\} \cup \{(\alpha, n) : n \geq \varphi(\alpha)\}$  where  $\varphi \in \omega^\kappa$ .

For a topological space  $X$ , the *tightness* of  $X$ ,  $t(X)$ , is the smallest cardinal  $\lambda$  such that for every point  $x \in X$  and  $A \subseteq X$ , if  $x \in \text{cl } A$  then there exists  $B \subseteq A$  with  $|B| \leq \lambda$  and  $x \in \text{cl } B$ .

It is easily seen that  $t(S_\kappa) = \omega$  for each  $\kappa$ . But the tightness of the product space of two sequential fans,  $t(S_\omega \times S_\kappa)$ , is more complicated. In the paper [5] we define a

\* Correspondence address: Masaru Kada, University of Osaka Prefecture, College of Integrated Arts and Sciences (graduate course), 1-1, Gakuen-Cho, Sakai-City, Osaka 593, Japan.

E-mail: kada@center.osakafu-u.ac.jp.

<sup>1</sup> E-mail: yuasa@logic.info.waseda.ac.jp.

new cardinal invariant  $\mathfrak{b}^*$  and showed that it gives a combinatorial characterization of the tightness of  $S_\omega \times S_\kappa$ .

For  $f, g \in \omega^\omega$ ,  $f \leq^* g$  if  $f(n) \leq g(n)$  for all but finitely many  $n \in \omega$ .  $\mathfrak{b}$  is the smallest size of the unbounded family of  $\omega^\omega$  with respect to  $\leq^*$ , and  $\mathfrak{d}$  is the smallest size of the dominating family of  $\omega^\omega$  with respect to  $\leq^*$ . Clearly  $\mathfrak{b} \leq \mathfrak{d}$  holds.

We define the cardinal  $\mathfrak{b}^*$  and state our results proved in [5].

**Definition 1.1.**  $\mathfrak{b}^*$  is the smallest cardinal  $\lambda$  such that, for every unbounded family  $\mathcal{F} \subseteq \omega^\omega$ , there exists a subfamily  $\mathcal{G} \subseteq \mathcal{F}$  such that  $|\mathcal{G}| \leq \lambda$  and  $\mathcal{G}$  is still unbounded.

**Theorem 1.2** [5, Theorem 1.3].

- (1) For  $\omega \leq \kappa < \mathfrak{b}$ ,  $t(S_\omega \times S_\kappa) = \omega$  holds.
  - (2)  $t(S_\omega \times S_\mathfrak{b}) = \mathfrak{b}$ .
  - (3) For  $\kappa \geq \mathfrak{b}^*$ ,  $t(S_\omega \times S_\kappa) = \mathfrak{b}^*$  holds.
- In particular,  $t(S_\omega \times S_{2^\omega}) = \mathfrak{b}^*$ .

**Theorem 1.3** [5, Theorem 1.4].

- (1)  $\mathfrak{b} \leq \mathfrak{b}^* \leq \mathfrak{d}$ .
- (2) Both  $\mathfrak{b} < \mathfrak{b}^*$  and  $\mathfrak{b}^* < \mathfrak{d}$  are consistent with ZFC.

Now we are interested in the cardinal  $\mathfrak{b}^*$  itself rather than sequential fans. In the present paper we introduce some cardinal invariants which generalize the notion of  $\mathfrak{b}^*$  and investigate their properties.

Let  $\mathcal{I}$  denote a  $\sigma$ -ideal on  $\omega^\omega$  containing all singletons. The cardinal  $\text{non}(\mathcal{I})$  is the smallest size of a subset of  $\omega^\omega$  which is not in  $\mathcal{I}$ , and the cardinal  $\text{cof}(\mathcal{I})$  is the smallest size of a basis of the ideal  $\mathcal{I}$ , that is, the smallest size of a subfamily  $\mathcal{A}$  of  $\mathcal{I}$  such that for every  $B \in \mathcal{I}$  there exists  $A \in \mathcal{A}$  with  $B \subseteq A$ . It is easily seen that  $\text{non}(\mathcal{I}) \leq \text{cof}(\mathcal{I})$  holds for each ideal  $\mathcal{I}$ .

Analogously to Definition 1.1, we define the following cardinal for the ideal  $\mathcal{I}$ .

**Definition 1.4.**  $\text{shr}(\mathcal{I})$  is the smallest cardinal  $\lambda$  such that, for every set  $A \subseteq \omega^\omega$  with  $A \notin \mathcal{I}$ , there exists a subset  $B \subseteq A$  such that  $|B| \leq \lambda$  and  $B \notin \mathcal{I}$ .

Let  $\mathcal{N}$ ,  $\mathcal{M}$ ,  $\mathcal{K}$  denote the ideals generated by null sets, meager sets, and  $\sigma$ -compact sets in the Baire space  $\omega^\omega$  respectively. Note that  $\mathcal{K}$  is a subideal of  $\mathcal{M}$ , and it is not so hard to see that  $\mathfrak{b} = \text{non}(\mathcal{K})$ ,  $\mathfrak{d} = \text{cof}(\mathcal{K})$  and  $\mathfrak{b}^* = \text{shr}(\mathcal{K})$ .

For the cardinals  $\text{non}(\mathcal{M})$ ,  $\text{non}(\mathcal{N})$ ,  $\text{cof}(\mathcal{M})$ ,  $\text{cof}(\mathcal{N})$ ,  $\mathfrak{b}$  and  $\mathfrak{d}$ , the following inequalities are known. These inequalities form a part of *Cichoń's diagram*. (See [3,6] for more information.)

**Theorem 1.5.**

- (1)  $\mathfrak{b} \leq \text{non}(\mathcal{M})$ .
- (2) (Miller [12, Theorem 3])  $\mathfrak{d} \leq \text{cof}(\mathcal{M})$ .
- (3) (Bartoszyński [1])  $\text{cof}(\mathcal{M}) \leq \text{cof}(\mathcal{N})$ .

We shall prove the following relations, which generalize Theorem 1.3.

**Theorem 1.6.**

- (1)  $\text{non}(\mathcal{M}) \leq \text{shr}(\mathcal{M}) \leq \text{cof}(\mathcal{M})$ ,  $\text{non}(\mathcal{N}) \leq \text{shr}(\mathcal{N}) \leq \text{cof}(\mathcal{N})$ .
- (2) Each of the following sentences is consistent with ZFC:  $\text{non}(\mathcal{M}) < \text{shr}(\mathcal{M})$ ,  $\text{shr}(\mathcal{M}) < \text{cof}(\mathcal{M})$ ,  $\text{non}(\mathcal{N}) < \text{shr}(\mathcal{N})$ ,  $\text{shr}(\mathcal{N}) < \text{cof}(\mathcal{N})$ .

Our notation is standard and we refer the reader to [2] or [10] for undefined notions.

## 2. Consistency results

In this section we show that the cardinals  $\text{shr}(\mathcal{M})$  and  $\text{shr}(\mathcal{N})$  are nontrivial, that is, they are consistently different from the cardinals which appear in Cichon's diagram.

**Theorem 2.1.** *For any ideal  $\mathcal{I}$  on  $\omega^\omega$ ,  $\text{non}(\mathcal{I}) \leq \text{shr}(\mathcal{I}) \leq \text{cof}(\mathcal{I})$  holds.*

**Proof.**  $\text{non}(\mathcal{I}) \leq \text{shr}(\mathcal{I})$  follows immediately from the definition of  $\text{shr}(\mathcal{I})$ . To show  $\text{shr}(\mathcal{I}) \leq \text{cof}(\mathcal{I})$ , fix  $X \subseteq \omega^\omega$  with  $X \notin \mathcal{I}$  and let  $\mathcal{A} = \{A_\xi: \xi < \text{cof}(\mathcal{I})\}$  a basis of the ideal  $\mathcal{I}$ . For each  $\xi < \text{cof}(\mathcal{I})$  we can find  $y_\xi \in X \setminus A_\xi$ . Let  $Y = \{y_\xi: \xi < \text{cof}(\mathcal{I})\} \subseteq X$ . Then  $|Y| \leq \text{cof}(\mathcal{I})$  and  $Y \notin \mathcal{I}$ .  $\square$

Theorem 1.6(1) follows immediately from the above theorem.

Now we turn to the consistency proofs. All required models are obtained by Cohen or random forcing extensions. Through this section  $M$  is a c.t.m. for ZFC,  $\mathbb{P}$  is a forcing notion satisfying c.c.c., and  $G$  is a  $\mathbb{P}$ -generic filter over  $M$ .

Note that every Borel subset of  $\omega^\omega$  can be coded by a function in  $\omega^\omega$ , which is called a *Borel code*. (See [8,11] for details.) For a Borel code  $c$ , let  $\widehat{c}$  denote the Borel set coded by  $c$ .

It is known that the ideals  $\mathcal{M}$ ,  $\mathcal{N}$  and  $\mathcal{K}$  are generated by Borel sets and absolute in the sense of Borel codes, that is, for every Borel code  $c$  the sentence “ $\widehat{c}$  is meager” (or “ $\widehat{c}$  is measure zero” or “ $\widehat{c}$  is  $\sigma$ -compact” respectively) is absolute for all transitive models of ZFC containing  $c$  as a Borel code [11].

**Definition 2.2.** Let  $\mathcal{I}$  denote a  $\sigma$ -ideal on  $\omega^\omega$  which is absolute in the sense of Borel codes. We say  $\mathbb{P}$  *preserves  $\mathcal{I}$ -positive sets* if for any  $A \subseteq \omega^\omega$  with  $A \notin \mathcal{I}$  we have  $\Vdash_{\mathbb{P}} A \notin \mathcal{I}$ .

Needless to say, “ $\mathcal{M}$ -positive” means “nonmeager”, “ $\mathcal{N}$ -positive” means “outer measure positive”, and “ $\mathcal{K}$ -positive” means “unbounded”.

For an infinite set  $I$ , let  $\mathbb{C}(I)$ ,  $\mathbb{B}(I)$  denote the Cohen and random forcing notions with the index set  $I$  respectively (see [2,10,11]).

**Lemma 2.3** [4, Corollary 3.5]. *For any infinite set  $I$ ,  $\mathbb{C}(I)$  preserves unbounded sets.*

**Lemma 2.4** [2, Subsections 6.2.2 and 6.2.3]. *For any infinite set  $I$ ,  $\mathbb{C}(I)$  preserves non-meager sets, and  $\mathbb{B}(I)$  preserves sets having outer measure positive.*

**Definition 2.5.** For a forcing notion  $\mathbb{P}$ , a standard  $\mathbb{P}$ -name  $\dot{f}$  for a real is a name uniquely determined by a system  $\{A_{mn}: m, n < \omega\}$  with the following:

- (1)  $A_{mn} \subseteq \mathbb{P}$  is an antichain of  $\mathbb{P}$  and  $n \neq n'$  implies  $A_{mn} \cap A_{mn'} = \emptyset$ ,
- (2)  $\bigcup_{n < \omega} A_{mn}$  is a maximal antichain of  $\mathbb{P}$ , and
- (3) For each  $p \in A_{mn}$ ,  $p \Vdash_{\mathbb{P}} \dot{f}(m) = n$ .

**Theorem 2.6.** *Assume  $2^\omega = \lambda$  and  $\kappa$  be an infinite cardinal. Then,*

- (1) *for  $\mathbb{P} = \mathbb{C}(\kappa)$  and  $\mathcal{I} = \mathcal{K}$  or  $\mathcal{I} = \mathcal{M}$ , or*
- (2) *for  $\mathbb{P} = \mathbb{B}(\kappa)$  and  $\mathcal{I} = \mathcal{N}$ ,*

*the following holds: any subset  $A$  of  $\omega^\omega$  with  $A \notin \mathcal{I}$  has a subset  $B \subseteq A$  such that  $|B| \leq \lambda$  and  $B \notin \mathcal{I}$  in the forcing extension by  $\mathbb{P}$ .*

**Proof.** We prove both in the cases (1) and (2) in parallel. Let  $\mathbb{P}(I)$  denote  $\mathbb{C}(I)$  when  $\mathbb{P} = \mathbb{C}(\kappa)$  and  $\mathbb{B}(I)$  when  $\mathbb{P} = \mathbb{B}(\kappa)$ . Note that for every condition  $p \in \mathbb{P}$  we can find a countable  $J \subseteq \kappa$  and  $p' \in \mathbb{P}(J)$  so that  $p = p' \times 1_{\mathbb{P}(\kappa \setminus J)}$ . We denote such a countable set  $J$  by  $\text{supp}(p)$ .

For an infinite  $I \subseteq \kappa$ , let  $X(I)$  be the collection of all standard  $\mathbb{P}(I)$ -names for reals and let  $\mathcal{X} = X(\kappa)$ . It suffices to deal with the case  $\kappa > \lambda$ .

Suppose the theorem is false, then there are  $p_0 \in \mathbb{P}(\kappa)$  and a collection  $\dot{A}$  of standard  $\mathbb{P}(\kappa)$ -names for reals such that

$$p_0 \Vdash_{\mathbb{P}(\kappa)} "\dot{A} \notin \mathcal{I} \wedge \forall B \subseteq \dot{A} (|B| \leq \lambda \rightarrow B \in \mathcal{I})".$$

Let  $S = \{X(I): I \in [\kappa]^\lambda \wedge \text{supp}(p_0) \subseteq I\}$ , then  $S \subseteq [\mathcal{X}]^\lambda$ .  $S$  is stationary, since it is unbounded and closed under unions of increasing  $\omega_1$ -sequences. By assumption and the fact that  $\mathbb{P}(I)$  preserves  $\mathcal{I}$ -positive sets for any infinite set  $I$ , for each  $X = X(I) \in S$  we get a standard  $\mathbb{P}(I)$ -name  $\dot{c}_X$  for a Borel code so that

$$p_0 \Vdash_{\mathbb{P}(I)} "\dot{A} \cap X \subseteq \widehat{\dot{c}_X} \wedge \widehat{\dot{c}_X} \in \mathcal{I}."$$

Note that  $\dot{c}_X \in X = X(I)$ . By Fodor's lemma for  $[\mathcal{X}]^\lambda$  (see [7, Theorem 3.2]) there is a stationary set  $S' \subseteq S$  and a standard  $\mathbb{P}(\kappa)$ -name  $\dot{c}$  for a Borel code such that  $\dot{c}_X = \dot{c}$  for all  $X \in S'$ . Clearly  $p_0 \Vdash_{\mathbb{P}(\kappa)} "\widehat{\dot{c}} \in \mathcal{I}"$ . Since  $S'$  is unbounded in  $[\mathcal{X}]^\lambda$ , we have  $p_0 \Vdash_{\mathbb{P}(\kappa)} "\dot{x} \in \widehat{\dot{c}}"$  for all  $\dot{x} \in \dot{A}$ . It means that  $p_0 \Vdash_{\mathbb{P}(\kappa)} "\dot{A} \subseteq \widehat{\dot{c}}"$  and hence  $p_0 \Vdash_{\mathbb{P}(\kappa)} "\dot{A} \in \mathcal{I}"$ , which is a contradiction.  $\square$

**Corollary 2.7.** *Assume CH and let  $\kappa$  be a cardinal of uncountable cofinality. Then,*

- (1)  *$\mathfrak{b} = \mathfrak{b}^* = \text{non}(\mathcal{M}) = \text{shr}(\mathcal{M}) = \omega_1$  and  $\mathfrak{d} = \text{cof}(\mathcal{M}) = \kappa$  hold in the forcing model by  $\mathbb{C}(\kappa)$ .*
- (2)  *$\text{non}(\mathcal{N}) = \text{shr}(\mathcal{N}) = \omega_1$  and  $\text{cof}(\mathcal{N}) = \kappa$  hold in the forcing model by  $\mathbb{B}(\kappa)$ .*

Using Lemmas 2.3, 2.4 and Theorem 2.6, we can easily prove all the remaining consistency results.

**Lemma 2.8.** *Let  $\mathcal{I}$  denote a  $\sigma$ -ideal on  $\omega^\omega$  which is absolute in the sense of Borel codes. If  $\mathbb{P}$  preserves  $\mathcal{I}$ -positive sets, then  $(\text{shr}(\mathcal{I}))^M \leq (\text{shr}(\mathcal{I}))^{M[G]}$  holds.*

**Proof.** Fix  $\lambda < \text{shr}(\mathcal{I})$  arbitrarily. Let  $A \subseteq \omega^\omega \cap M$  be the set such that  $A \notin \mathcal{I}$  and for  $B \subseteq A$  if  $|B| \leq \lambda$  then  $B \in \mathcal{I}$ . We show that  $A$  witnesses  $(\text{shr}(\mathcal{I}))^{M[G]} > \lambda$ .

Since  $\mathbb{P}$  preserves  $\mathcal{I}$ -positive sets,  $\Vdash_{\mathbb{P}} "A \notin \mathcal{I}"$  also holds. Fix a  $\mathbb{P}$ -name  $\dot{B}$  such that  $\Vdash_{\mathbb{P}} "\dot{B} \subseteq A \wedge |\dot{B}| \leq \lambda"$ . Since  $\mathbb{P}$  is c.c.c., we can find  $B' \subseteq A$  such that  $|B'| \leq \lambda$  and  $\Vdash_{\mathbb{P}} "\dot{B} \subseteq B'"$ . By assumption, we can find a Borel code  $c$  so that  $B' \subseteq \hat{c} \in \mathcal{I}$  holds. By the absoluteness of  $\mathcal{I}$ ,

$$\Vdash_{\mathbb{P}} "\dot{B} \subseteq B' \subseteq (\hat{c})^M = \hat{c} \cap M \subseteq \hat{c} \in \mathcal{I}",$$

so  $\Vdash_{\mathbb{P}} "\dot{B} \in \mathcal{I}"$  holds.  $\square$

**Corollary 2.9.** *Assume  $\text{MA} + \omega_1 < 2^\omega = \lambda \leq \kappa$  and  $\kappa$  has uncountable cofinality. Then,*

- (1)  $\mathfrak{b} = \text{non}(\mathcal{M}) = \omega_1$ ,  $\mathfrak{b}^* = \text{shr}(\mathcal{M}) = \lambda$  and  $\mathfrak{d} = \text{cof}(\mathcal{M}) = \kappa$  hold in the forcing model by  $\mathbb{C}(\kappa)$ .
- (2)  $\text{non}(\mathcal{N}) = \omega_1$ ,  $\text{shr}(\mathcal{N}) = \lambda$  and  $\text{cof}(\mathcal{N}) = \kappa$  hold in the forcing model by  $\mathbb{B}(\kappa)$ .

### 3. Adding a Cohen real can change $\mathfrak{b}^*$

We notice the fact that  $(\mathfrak{b}^*)^M = (\mathfrak{b}^*)^{M[G]}$  holds in all the forcing models obtained in Corollaries 2.7 and 2.9. So it is natural to ask whether  $\mathfrak{b}^*$  is absolute for Cohen or random forcing extensions. In fact, the answer is ‘no’ for Cohen forcing. This section is devoted to show that adding a Cohen real can change  $\mathfrak{b}^*$ .

Here we consider meager or nonmeager sets in the Cantor space  $2^\omega$  instead of the Baire space  $\omega^\omega$ . The cardinals associated with the meager ideal are the same if we used either of two spaces.

For  $x, y \in 2^\omega$ ,  $x + y$  is defined as  $(x + y)(n) = x(n) + y(n) \bmod 2$  for each  $n \in \omega$ . Similarly, for  $s, t \in 2^{<\omega}$ ,  $s + t$  is defined as  $\text{dom}(s + t) = \min\{|s|, |t|\}$  and  $(s + t)(n) = s(n) + t(n) \bmod 2$  for  $n \in \text{dom}(s + t)$ . For  $A \subseteq 2^\omega$  and  $z \in 2^\omega$  we define  $A + z = \{x + z: x \in A\}$ .

For  $r \in [\omega]^\omega$  the increasing enumeration of  $r$  is denoted by  $f_r$ . Then we will say  $X \subseteq [\omega]^\omega$  is *bounded* if the family of functions  $\{f_r: r \in X\}$  is bounded with respect to  $\leq^*$ , and otherwise we say it is *unbounded*. In the following argument we will identify an infinite subset of  $\omega$  with its characteristic function. It is easy to see that every nonmeager set in  $2^\omega$  is unbounded.

**Theorem 3.1.** *Let  $\kappa$  be a cardinal and  $X \in M$  be a nonmeager subset of  $2^\omega$  such that, for  $Y \in M$  if  $Y \subseteq X$  and  $|Y| < \kappa$  then  $Y$  is meager. Let  $c$  be a Cohen real over  $M$ . Then in  $M[c]$ ,*

- (1)  $X + c$  is unbounded and
- (2) Every  $Z \subseteq X + c$  of size less than  $\kappa$  is bounded.

Because any element of  $X + c$  takes infinitely often the value 1, (1) and (2) make sense.

**Corollary 3.2.**  $(\text{shr}(\mathcal{M}))^M \leq (\mathfrak{b}^*)^{M[c]}$ . In particular, if  $(\mathfrak{b}^*)^M < (\text{shr}(\mathcal{M}))^M$ , then  $(\mathfrak{b}^*)^M < (\mathfrak{b}^*)^{M[c]}$ .

Of course,  $\mathfrak{b}^* < \text{shr}(\mathcal{M})$  is consistent with ZFC. (Consider the model obtained by adding  $\omega_2$  random reals to the model of CH.)

A subtree  $T$  of  $2^{<\omega}$  with no maximal node is called *nowhere dense* if for any  $t \in 2^{<\omega}$  there exists  $s \in 2^{<\omega} \setminus T$  such that  $t \subseteq s$ . Note that for any  $X \subseteq 2^\omega$  the subtree  $T(X) = \{r \upharpoonright n : r \in X \wedge n \in \omega\}$  of  $2^{<\omega}$  is nowhere dense if and only if  $X$  is nowhere dense in  $2^\omega$ .

**Lemma 3.3.** Let  $T$  be a nowhere dense tree and  $k \in \omega$ . Then for any  $p \in 2^{<\omega}$  there exists  $q \in 2^{<\omega}$  with  $p \subseteq q$  such that:

$$\forall t \in \text{Lev}_{|q|}(T) \ (k < |(t+q)^{-1}\{1\}|)$$

**Proof.** Fix a nowhere dense tree  $T$  and  $p \in 2^{<\omega}$  arbitrarily. For  $q \in 2^{<\omega}$  with  $p \subseteq q$  and  $t \in \text{Lev}_{|q|}(T)$  we say  $t$  survives  $q$  if

$$|(t+p)^{-1}\{1\}| = |(t+q)^{-1}\{1\}|.$$

In order to prove the lemma it suffices to show that there exists  $q \in 2^{<\omega}$  with  $p \subseteq q$  such that no node in  $\text{Lev}_{|q|}(T)$  survives  $q$ . So we will inductively construct a finite sequence  $p = p_0 \subset p_1 \subset \dots \subset p_l$  such that:

- $|\{t \in \text{Lev}_{|p_i|}(T) : t \text{ survives } p_i\}| > |\{t \in \text{Lev}_{|p_{i+1}|}(T) : t \text{ survives } p_{i+1}\}|$  for  $i < l$  and
- $|\{t \in \text{Lev}_{|p_l|}(T) : t \text{ survives } p_l\}| = 0$ ,

and let  $q = p_l$ . Assume that  $p_i$  is constructed. If there is no node  $t \in \text{Lev}_{|p_i|}(T)$  surviving  $p_i$ , let  $l = i$  and we are done. Otherwise, pick a node  $t \in \text{Lev}_{|p_i|}(T)$  surviving  $p_i$ . Since  $T$  is nowhere dense, there exists  $s \in T$  such that  $t \subset s$  and  $s \in 2^{<\omega} \setminus T$ . We define  $p_{i+1} \in 2^{|s|}$  as follows.

$$p_{i+1}(n) = \begin{cases} p_i(n) & (n < |p_i|), \\ s(n) & (|p_i| \leq n < |s|). \end{cases}$$

Since  $t \in \text{Lev}_{|p_{i+1}|}(T)$  survives  $p_{i+1}$  if and only if  $t$  and  $p_{i+1}$  coincide on  $|p_{i+1}| \setminus |p_i|$  and  $t \upharpoonright |p_i|$  survives  $p_i$ , it is clear that this  $p_{i+1}$  satisfies the condition.  $\square$

**Proof of Theorem 3.1.** (1) It is clear from the fact mentioned above, because  $X$  is still nonmeager in  $M[c]$  by Lemma 2.3 and hence so is  $X + c$ .

(2) There is  $Y \in M$  of the same size as  $Z$  such that  $Y \subseteq X$  and  $Z \subseteq Y + c$ . Then we will work in  $M$ . Since  $Y$  is meager, there is a countable family  $\{Y_m : m \in \omega\}$  of nowhere dense sets which covers  $Y$ . Let  $T_m$  be the nowhere dense tree  $T(Y_m)$ . Fix  $m \in \omega$  and we will construct a  $\mathbb{C}$ -name  $\dot{g}_m$  such that for any infinite path  $r$  through  $T_m$  we have

$\Vdash \forall n < \omega (f_{r+c}(n) \leq \dot{g}_m(n))$ . We will use  $2^{<\omega}$  with inverse inclusion  $\supseteq$  as our Cohen forcing notion  $\mathbb{C}$  instead of  $\text{Fn}(\omega, 2)$ . Using Lemma 3.3 we get a maximal antichain  $A_k$  for each  $k \in \omega$  so that  $k < |(t+q)^{-1}\{1\}|$  for any  $q \in A_k$  and any  $t \in \text{Lev}_{|q|}(T)$ . We define a  $\mathbb{C}$ -name  $\dot{g}_m$  as

$$\Vdash \dot{g}_m(k) = |q| \text{ iff } \dot{G} \cap A_k = \{q\}$$

for any  $k \in \omega$ . Clearly this  $\dot{g}_m$  is what we need. At last we take a  $\mathbb{C}$ -name  $\dot{g}$  so that  $\Vdash_{\mathbb{C}} \text{"}\forall m \in \omega (\dot{g}_m \leq^* \dot{g})\text{"}$ , then it witnesses the boundedness of  $Z$ .  $\square$

#### 4. Cofinality of $\mathfrak{b}^*$

It is well known that  $\mathfrak{b}$  is regular and  $\mathfrak{d}$  has uncountable cofinality. (More precisely,  $\text{cf}(\mathfrak{d}) \geq \mathfrak{b}$ .) Now we observe that  $\mathfrak{b}^*$  also has uncountable cofinality.

**Definition 4.1.** For  $A, B \subseteq \omega$ ,  $A \subseteq^* B$  if  $A \setminus B$  is finite. A family  $\{B_\alpha: \alpha < \lambda\}$  of infinite subsets of  $\omega$  is called a *tower* if  $B_\beta \subseteq^* B_\alpha$  for  $\alpha < \beta < \lambda$  and there is no infinite  $B \subseteq \omega$  satisfying  $B \subseteq^* B_\alpha$  for all  $\alpha < \lambda$ .

**Definition 4.2.**  $\mathfrak{t}$  is the smallest size of a tower.

It is known that  $\mathfrak{t}$  is regular and  $\omega_1 \leq \mathfrak{t} \leq \mathfrak{b}$  (see [13, Theorem 3.1]). The following theorem shows that  $\mathfrak{b}^*$  has uncountable cofinality.

**Theorem 4.3.**  $\text{cf}(\mathfrak{b}^*) \geq \mathfrak{t}$ .

**Proof.** Assume that  $\text{cf}(\mathfrak{b}^*) = \lambda < \mathfrak{t}$ . Then there are  $\kappa_\alpha, \mathcal{F}_\alpha$  for  $\alpha < \lambda$  such that

- (1)  $\kappa_\alpha < \kappa_\beta < \mathfrak{b}^*$  for  $\alpha < \beta < \lambda$  and  $\sup\{\kappa_\alpha: \alpha < \lambda\} = \mathfrak{b}^*$ ,
- (2)  $\mathcal{F}_\alpha$  is an unbounded family of non-decreasing functions in  $\omega^\omega$  and  $|\mathcal{F}_\alpha| = \kappa_\alpha$ ,
- (3) For any  $\mathcal{F} \subseteq \mathcal{F}_\alpha$  if  $|\mathcal{F}| < \kappa_\alpha$  then  $\mathcal{F}$  is bounded.

For each  $\alpha < \lambda$ , let  $\mathcal{S}_\alpha = [\mathcal{F}_\alpha]^{<\kappa_\alpha}$ . Set

$$\mathcal{S} = \left\{ S \in \prod_{\alpha < \lambda} \mathcal{S}_\alpha: \exists \beta < \lambda \forall \alpha < \lambda (|S(\alpha)| \leq \kappa_\beta) \right\}.$$

Then for any  $S \in \mathcal{S}$ ,  $\bigcup_{\alpha < \lambda} S(\alpha)$  is bounded. So we can take  $h_S \in \omega^\omega$  for each  $S \in \mathcal{S}$  such that  $f \leq^* h_S$  for all  $f \in \bigcup_{\alpha < \lambda} S(\alpha)$ .

Define a quasi-order  $\prec$  on  $\mathcal{S}$  by  $S \prec T$  iff there exists  $\beta < \lambda$  such that  $S(\alpha) \subseteq T(\alpha)$  for all  $\alpha$  with  $\beta \leq \alpha < \lambda$ . For each  $S \in \mathcal{S}$ , let  $\mathcal{H}_S = \{f \in \omega^\omega: \forall T \succ S (f \leq^* h_T)\}$ , and let  $\mathcal{H} = \bigcup_{S \in \mathcal{S}} \mathcal{H}_S$ . We shall show that  $\mathcal{H}$  is unbounded and that any subfamily  $\mathcal{G} \subseteq \mathcal{H}$  with  $|\mathcal{G}| < \mathfrak{b}^*$  is bounded, which contradicts  $\text{cf}(\mathfrak{b}^*) < \mathfrak{t} \leq \mathfrak{b}$ .

First we show that  $\mathcal{H}$  is unbounded. So take  $h \in \omega^\omega$  arbitrarily. By the assumption that  $\lambda < \mathfrak{t}$ , we can take  $f_\alpha \in \mathcal{F}_\alpha$  and infinite  $B_\alpha \subseteq \omega$  by induction on  $\alpha < \lambda$  satisfying the following:  $B_\alpha = \{k < \omega: h(k) < f_\alpha(k)\}$  for each  $\alpha < \lambda$ , and  $B_\beta \subseteq^* B_\alpha$  for  $\alpha < \beta < \lambda$ . Choose infinite  $B \subseteq \omega$  so that  $B \subseteq^* B_\alpha$  for all  $\alpha < \lambda$ .

Let  $S = \{\{f_\alpha\}: \alpha < \lambda\}$ , and define  $f \in \omega^\omega$  by  $f(k) = h(k) + 1$  for  $k \in B$  and  $f(k) = 0$  otherwise. Then  $f \not\leq^* h$ . On the other hand,  $f \leq f_\alpha$  for all  $\alpha < \lambda$ , and if  $T \succ S$  then there is  $\alpha < \lambda$  such that  $f_\alpha \leq^* h_T$ , so  $f \in \mathcal{H}_S$  and hence  $f \in \mathcal{H}$ .

Next we show that  $\mathcal{G}$  is bounded for all  $\mathcal{G} \subseteq \mathcal{H}$  with  $|\mathcal{G}| < \mathfrak{b}^*$ . Fix such  $\mathcal{G} \subseteq \mathcal{H}$  and take  $\beta_0 < \lambda$  so that  $|\mathcal{G}| < \kappa_{\beta_0}$ . For each  $g \in \mathcal{G}$ , take  $S_g \in \mathcal{S}$  and  $\beta_g < \lambda$  such that  $g \in \mathcal{H}_{S_g}$  and  $|S_g(\alpha)| \leq \kappa_{\beta_g}$  for all  $\alpha < \lambda$ . For each  $\alpha \in (\beta_0, \lambda)$ , let  $\mathcal{G}_\alpha = \{g \in \mathcal{G}: \beta_g < \alpha\}$ . We claim that  $\mathcal{G}_\alpha$  is bounded for all  $\alpha \in (\beta_0, \lambda)$ , which implies that  $\mathcal{G} = \bigcup_{\beta_0 < \alpha < \lambda} \mathcal{G}_\alpha$  is bounded. So fix  $\alpha \in (\beta_0, \lambda)$ . Since  $\bigcup_{g \in \mathcal{G}_\alpha} S_g(\xi) \in \mathcal{S}_\xi$  for all  $\xi \in (\alpha, \lambda)$ , we can define  $T \in \mathcal{S}$  by  $T(\xi) = \bigcup_{g \in \mathcal{G}_\alpha} S_g(\xi)$  if  $\alpha \leq \xi < \lambda$  and  $T(\xi) = \emptyset$  otherwise. Since  $g \in S_g$  and  $S_g \prec T$  for all  $g \in \mathcal{G}_\alpha$ , we have  $g \leq^* h_T$  for all  $g \in \mathcal{G}_\alpha$ .  $\square$

## 5. Questions

As we mentioned in Section 1,  $\mathfrak{b} \leq \text{non}(\mathcal{M})$  and  $\mathfrak{d} \leq \text{cof}(\mathcal{M})$  hold in ZFC. So it is natural to ask the following question.

**Question 1.** Does  $\mathfrak{b}^* \leq \text{shr}(\mathcal{M})$  hold in ZFC?

In Section 3 we proved that adding a Cohen real can change  $\mathfrak{b}^*$ .

**Question 2.** Can adding random reals change  $\mathfrak{b}^*$ ?

Note that the dual result to Corollary 3.2 cannot hold, because adding random reals cannot raise  $\mathfrak{d}$ , and  $\mathfrak{d} < \text{non}(\mathcal{N})$  is consistent with ZFC (see [3]).

**Question 3.** Is  $(\text{shr}(\mathcal{M}))^M = (\mathfrak{b}^*)^{M[G]}$  for Cohen forcing?

**Question 4.** What happens about  $\text{shr}(\mathcal{M})$  or  $\text{shr}(\mathcal{N})$  when Cohen or random reals are added?

In Section 4 we proved that  $\text{cf}(\mathfrak{b}^*) \geq \mathfrak{t}$ . But it is known that  $\mathfrak{t} < \mathfrak{b}$  is consistent with ZFC (see [13, Theorem 5.3]).

**Question 5.** Does  $\text{cf}(\mathfrak{b}^*) \geq \mathfrak{b}$  hold in ZFC?

*Added in proof.* After the submission of this paper, the second author proved that  $\mathfrak{b}^* \leq \text{shr}(\mathcal{M})$ , which gives the answer to Question 1. On the other hand, the first author proved that  $\text{shr}(\mathcal{N}) < \mathfrak{b}$  is consistent with ZFC. So we have proved all ZFC-provable inequalities between two cardinals from this paper. We will publish these results in the near future.



## Acknowledgements

We thank Prof. K. Eda and S. Kamo for giving helpful advice for this research. The result in Section 4 is due to Kamo and we thank him for permitting us to include it to this paper. We also thank S. Taniyama and H. Fujita for suggesting the idea of the notions of  $\mathfrak{b}^*$ ,  $\text{shr}(\mathcal{M})$  and  $\text{shr}(\mathcal{N})$ .

## References

- [1] T. Bartoszyński, Combinatorial aspects of measure and category, *Fund. Math.* 127 (1987) 225–239.
- [2] T. Bartoszyński and H. Judah, *Set Theory: On the Structure of the Real Line* (A.K. Peters, 1995).
- [3] T. Bartoszyński, H. Judah and S. Shelah, The Cichoń diagram, *J. Symbolic Logic* 58 (1993) 401–423.
- [4] J. Cichoń and J. Pawlikowski, On ideals of subsets of the plane and on Cohen reals, *J. Symbolic Logic* 51 (1986) 560–569.
- [5] K. Eda, M. Kada and Y. Yuasa, The tightness about sequential fans and combinatorial properties, to appear in *J. Math. Soc. Japan*.
- [6] D.H. Fremlin, Cichoń's diagram, in: G. Choquet, M. Rogalski and J. Saint-Raymond, eds., *Séminaire Initiation à l'Analyse* (Univ. Pierre et Marie Curie, 1983/84) (5–01)–(5–13).
- [7] T. Jech, Some combinatorial problems concerning uncountable cardinals, *Ann. Math. Logic* 5 (1973) 165–198.
- [8] T. Jech, *Set Theory* (Academic Press, 1978).
- [9] M. Kada, Tightness and combinatorial properties concerning sequential fans, Master's thesis (Univ. of Tsukuba, 1995).
- [10] K. Kunen, *Set Theory* (North-Holland, 1980).
- [11] K. Kunen, Random and Cohen reals, in: K. Kunen and J.E. Vaughan, eds., *Handbook of Set-Theoretic Topology* (North-Holland, Amsterdam, 1984) 887–911.
- [12] A.W. Miller, Additivity of measure implies dominating reals, *Proc. Amer. Math. Soc.* 91 (1984) 111–117.
- [13] E.K. van Douwen, The integers and topology, in: K. Kunen and J.E. Vaughan, eds., *Handbook of Set-Theoretic Topology* (North-Holland, Amsterdam, 1984) 111–167.